



UNIVERSITI PUTRA MALAYSIA

**AZOXYSTROBIN SENSITIVITY, MORPHOLOGICAL
AND GENETIC MARKER STUDIES OF
COLLETOTRICHUM GLOEOSPORIODES
(PENZ) PENZ. AND SACC. FROM MANGO AND
C CAPSICI (H. SYD.) E. BUTL AND BISBY FROM CHILLI**

QUAH JU LEE

FSAS 2000 22

**AZOXYSTROBIN SENSITIVITY, MORPHOLOGICAL
AND GENETIC MARKER STUDIES OF
COLLETOTRICHUM GLOEOSPORIOIDES '
(PENZ) PENZ. AND SACC. FROM MANGO AND
C. CAPSICI (H. SYD.) E. BUTL. AND BISBY FROM CHILLI**

By

QUAH JU LEE

**Thesis Submitted in Fulfilment of the Requirements for the Degree of
Master of Science in the Faculty of Science and Environmental Studies,
Universiti Putra Malaysia.**

December 2000



Abstract of thesis presented to the senate of Universiti Putra Malaysia in fulfilment
of the requirements for the degree of Masters of Science

**AZOXYSTROBIN SENSITIVITY, MORPHOLOGICAL AND GENETIC
MARKER STUDIES OF *COLLETOTRICHUM GLOEOSPORIOIDES* (PENZ.)
PENZ. AND SACC. FROM MANGO AND *C. CAPSICI* (H. SYD.) E. BUTL.
AND BISBY FROM CHILLI**

By

QUAH JU LEE

December 2000

Chairperson : Dr. Faridah Abdullah

Faculty : Science and Environmental Studies

This project evaluated the sensitivities of *Colletotrichum gloeosporioides* and *C. capsici* towards the fungicide Azoxystrobin. This was complemented by morphological and genetic marker studies of the two species in order to record the natural variation of the two species.

Twenty five cultures of *C. gloeosporioides* were isolated from mangoes (*Mangifera indica*) and 22 of cultures of *C. capsici* isolated from chillies (*Capsicum annum*) were used. They were collected from Malaysia, Indonesia, Thailand, Philippines and Taiwan. Single spore cultures were made to establish the species identity.

Isolates of both species were tested separately for fungicide sensitivity. The spore suspension was optimised to 10^5 spores/ml for bioassays. The assay medium used was a glycerol based Alkyl Ester broth, amended with rates of technical azoxystrobin at 0.0001, 0.001, 0.01 and 0.1ppm with blanks of Alkyl Ester broth as controls.

The tests were divided into statistical validation and actual baseline studies. For validation studies, 6 isolates in 6 replicate tests were used in the calculation of experimental errors. The baseline test consisted of 3 replicate treatments per Azoxystrobin rate per isolate with double controls for 14 isolates of *C. gloeosporioides* and 15 isolates of *C. capsici*. The validation isolates were included together with the baseline isolates during final analysis to give a total of 20 isolates for *C. gloeosporioides* and a total of 21 isolates for *C. capsici*.

The Alkyl Ester broth was inoculated with spore-suspensions of each isolate separately to give a final concentration of 10^5 spores/ml in the mixture. 60 μ l droplets of the broth-spore mixtures were incubated in 55 mm petri dishes and covered with cover slips. The reference isolate PML1 was included in all tests to monitor the consistency. Assessment was done by scoring the number of spores out of 50 that showed germ tube elongation to a minimum of twice the spore length.

Dose responses were plotted of log rate against the number of spores germinated to derive 80% effective dose or ED_{80} values for each isolate. This entire experiment was repeated twice and the data analysed to determine the intra and inter experimental variation. Sensitivity values were also calculated by linear regression analysis using arcsine transformation of the data (ACSAPWIN). Results showed that the ED_{80} range for *C. gloeosporioides* was between 0.01 ppm to 0.1 ppm while this was between 0.001 ppm to 0.01 ppm for *C. capsici*.

From cultural studies, *C. capsici* displayed similar characteristics whereas this range was more variable for *C. gloeosporioides*. Morphological studies showed that the growth rates of *C. gloeosporioides* and *C. capsici* were similar with an average growth rate of 7 cm/day for its diameter. The best growth medium was Alkyl Ester Agar followed by Oatmeal Agar, V8 Juice Agar, Potato Dextrose Agar, Malt Agar and lastly, Cooke's medium.

Genetic characterization was done using short primer Random Amplified Polymorphic DNA (RAPD) studies, Long primer-Random Amplified Polymorphic DNA studies (LP-RAPD) and Random Amplified Microsatellite studies (RAMS). The final dendrograms showed general clustering according to countries of origin for *C. gloeosporioides* and *C. capsici*.

Abstract testis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi syarat keperluan untuk ijazah Master Sains.

KAJIAN SENSITIVITI TERHADAP AZOXYSTROBIN DAN JUGA KAJIAN MORFOLOGI DAN 'MARKER' GENETIK UNTUK *COLLETOTRICHUM GLOEOSPORIODES* (PENZ) PENZ. AND SACC. DARI MANGA DAN *C. CAPSICI* (H. SYD.) E. BUTL. AND BISBY DARI CILI.

Oleh

QUAH JU LEE

Disember 2000

Penyelia : Dr. Faridah Abdullah

Fakulti : Sains dan Pengajian Alam Sekitar

Projek ini menilai sensitiviti *Colletotrichum gloeosporioides* dan *C. capsici* terhadap fungisid Azoxystrobin. Projek dilengkapi dengan kajian-kajian morfologi dan petanda genetik bagi kedua-dua species dengan tujuan untuk merekodkan maklumat variasi semulajadi bagi kedua-dua spesis.

Dua puluh lima kultur *C. gloeosporioides* yang dipencilkan dari buah mangga dan dua puluh dua kultur *C. capsici* dari buah cili telah digunakan. Kultur telah dikumpulkan dari Malaysia, Indonesia, Thailand, Filipina dan Taiwan. Sampel berpenyakit telah diisolatkan daripada perumahannya dan pemencilan dibuat untuk membolehkan pengenalpastian spesis dibuat seterusnya.

Isolat daripada spesis telah diuji secara berasingan untuk mendapatkan maklumat sensitivity terhadap fungisid. Suspensi isolat telah dioptimumkan pada kepekatan 10^5 spora/ml untuk ujian. Medium ujian yang telah digunakan adalah broth Alkyl Ester

yang berdasarkan gliserol dengan kepekatan Azoxystrobin pada 0.0001 ppm, 0.001 ppm, 0.01 ppm dan 0.1 ppm sementara kawalan yang digunakan adalah broth Alkyl Ester sahaja.

Ujian telah dibahagikan kepada kajian bacaan ralat ujikaji atau 'validation' dan kajian sebenar sensitivity atau 'baseline'. Untuk kajian 'validation'; 6 isolat telah digunakan dalam 6 ujian replikat dan data telah digunakan untuk mengira bacaan ralat eksperimen. Ujian 'baseline' terdiri daripada 3 rawatan replikat untuk setiap kadar bagi setiap isolat dengan dua kali ganda replikat masing-masing untuk rawatan kawalan bagi 14 isolat *C. gloeosporioides* dan 15 isolat *C. capsici*. Isolat dari kajian 'validation' kemudiannya dimasukkan bersama dengan isolat 'baseline' untuk memberikan sejumlah 20 isolat untuk *C. gloeosporioides* dan 21 isolat untuk *C. capsici*.

Broth Alkyl Ester telah diinokulat secara berasingan untuk ampaian spora setiap isolat untuk menghasilkan kepekatan akhir sebanyak 10^5 spora/ml dalam campuran tersebut. Titisan 60 μ l dari campuran broth dengan spora itu seterusnya dieramkan dalam piring petri 55 mm dengan ditutup dengan sisip kaca. Isolat rujukan, PML1, telah dimasukkan ke dalam semua ujian untuk memantau keseragaman ujian. Penilaian dilakukan dengan mengira bilangan spora yang menunjukkan percambahan melalui tiub germa sekurang-kurangnya dua kali ganda panjang spora tersebut dari 50 spora setiap plat.

'Dose responses' kemudiannya telah diplot bagi kadar log melawan bilangan spora tercampah untuk menghasilkan nilai-nilai ED_{50} untuk setiap isolat. Keseluruhan

ujian telah dijalankan dua kali untuk menentukan variasi sesama isolat dalam kumpulan atau 'intra' dan diantara isolat atau 'inter' dalam ujian-ujian. Nilai-nilai sensitiviti juga dikira melalui regresi linear dengan menggunakan transformasi arc-sin bagi data tersebut (ACSAPWIN). Keputusan menunjukkan nilai ED_{50} bagi *C. gloeosporioides* adalah diantara 0.01 ppm hingga 0.1 ppm sementara nilai ED_{50} bagi *C. capsici* adalah diantara 0.001 ppm hingga 0.01 ppm.

Dari kajian kultur *C. capsici* menunjukkan ciri-ciri yang hampir serupa sesama sendiri manakala ciri-ciri *C. gloeosporioides* menunjukkan lebih variasi. Kajian morfologi pula menunjukkan bahawa kadar pertumbuhan diameter hampir serupa dengan purata 7 cm/hari. Medium pertumbuhan yang paling baik adalah Agar Aitky Ester diikuti oleh Agar Oatmeal, Agar Jus, Agar Potato Dekrosa, Agar Malt Ektrak dan akhirnya Medium Cooke.

Pencirian genetik telah dilakukan dengan menggunakan kajian primer pendek RAPD, kajian primer panjang LP-RAPD dan kajian RAMS. Keputusan akhir dalam bentuk dendrogram menunjukkan perkelompokan umum mengikut negara asal masing-masing untuk *C. gloeosporioides* dan *C. capsici*.

ACKNOWLEDGEMENTS

My sincere gratitude to:

Zeneca Agrochemicals for sponsoring this project and especially to Steve Dale, Steve Heaney, Sally Davies and Maria Efstathiou of Zeneca Agrochemicals for all the technical advice given and kindness shown during my stay in Jealott's Hill, U.K.

All the staff of CCM Bioscience in Malacca for all their support and understanding.

Dr. Faridah Abdullah for her supervision of this project and for access to her facilities in the laboratory as well as her moral support during tough times.

Professor Tan Soon Guan for his supervision of the genetic studies which is included in Chapter 5 of the thesis.

Kam Boon Seng, my fiancée, for his moral support and kindness shown during the project.

Everyone in the Mycology and Genetic laboratories for their cheerful enthusiasm and understanding and help in times of need.

Especially to my beloved father, Dr. Quah Soon Cheang, for all his care, love and guidance.

Finally, to my beloved late mother who never failed to keep the faith and the belief that all things are possible with the help of God.

I certify that an Examination Committee met on 4th May 2001 to conduct the final examination of Quah Ju Lee on her Masters of Science thesis entitled "Azoxystrobin Sensitivity, Morphological and Genetic Marker Studies of *Colletotrichum gloeosporioides* (Penz) Penz. and Sacc. from Mango and *C. capsici* (H. Syd.) E. Butl. and Bisby from Chilli" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Ahlis of the Examination Committee are as follows:

Radzali Muse, Ph.D.
Associate Professor,
Faculty of Science and Environmental Studies,
Universiti Putra Malaysia.
(Chairman)

Faridah Abdullah, Ph.D.
Associate Professor,
Faculty of Science and Environmental Studies,
Universiti Putra Malaysia.
(Member)

Tan Soon Guan, Ph.D.
Professor,
Faculty of Science and Environmental Studies,
Universiti Putra Malaysia.
(Member)


Mah Shook Yin,
Research Officer,
Stesen Jalan Kebun,
MARDI, Klang,
Selangor.
(Member)



MOHD. GHAZALI MOHAYIDIN, Ph.D.,
Professor / Deputy Dean of Graduate School,
Universiti Putra Malaysia

Date: 27 JUN 2001

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted in fulfilment of the requirement for the degree of Masters of Science.



AINI IDERIS, Ph.D,
Professor,
Dean of Graduate School,
Universiti Putra Malaysia.

Date: 12 JUL 2001

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

Quah Ju Lee

TABLE OF CONTENTS

	PAGE
ABSTRACT	ii
ABSTRAK	v
ACKNOWLEDGEMENT	viii
APPROVAL SHEETS	ix
DECLARATION FORM	xi
LIST OF TABLES	xvi
LIST OF FIGURES	xvii
LIST OF ABBREVIATIONS	xx

CHAPTER

I	INTRODUCTION	1
II	LITERATURE REVIEW	
	Historical Background	6
	Synthesis of Azoxystrobin	8
	Mode of Action	9
	Biological Properties	10
	Physiological Properties	11
	Selection of <i>Colletotrichum</i> for Testing	12
	The Genus <i>Colletotrichum</i>	13
	<i>Colletotrichum gloeosporioides</i> (Penz.) Penz. and Sacc.	14
	<i>Colletotrichum capsici</i> (H. Syd.) E. Butl. And Bisby.	15
	Sample Storage	16
	Chilli	17
	Mango	18
	Resistance	19
	Measuring Resistance to Fungicides	21
	Baseline data assessment	22
	Bioassays for Detection	23
	Poison Plate Tests	25
	Liquid Broth Tests	25
	<i>In vivo</i> Tests	26
	Genetic Analysis	26
	Genetic Markers	27
	Various Genetic Methods	28
	Polymerase Chain Reaction	29
	Random Amplified Polymorphic DNA	30
	Long-Primer Random Amplified Polymorphic DNA	33
	Random Amplified Microsatellites	34

III	SAMPLE COLLECTION AND CHARACTERISATION	
	Introduction	36
	Objectives	37
	Material and Methods	38
	Collection of <i>C. gloeosporioides</i> and <i>C. capsici</i> isolates	38
	Isolation of Raw Samples	40
	Single Spore Isolation	40
	Species Identification	41
	Sample Storage	41
	Agar Slant Storage	41
	Cryopreservation	42
	Colony Characteristics	42
	Morphological and Growth Studies	43
	Sporulation, Spore Germination and Appresoria studies	44
	Pathogenicity Test	45
	Results	46
	Collection data for <i>Colletotrichum gloeosporioides</i> and <i>C. capsici</i>	46
	Culture Characteristics	49
	Solid Media Growth Studies	49
	Liquid Media Growth Studies	50
	Pathogenicity Studies	50
	Discussion	55
IV	FUNGICIDE SENSITIVITY BASELINE BIOASSAY	
	Introduction	59
	Material and Methods	60
	Media Used	60
	Optimization	61
	Spore Germination on Solid Media	61
	agar and broth	
	Washing of spores and increasing humidity on germination rates	64
	Using microcycle conidiation versus normal conidiation	64
	germination rates	
	Concentrations of inoculum and broth on germination rates	65
	Varying concentrations of Azoxystrobin on germination rates	65
	Droplet test development	65
	Validation bioassay	67
	Baseline bioassay	69
	Experimental controls	70
	Statistical Analysis	70
	Results	71
	The effect of different solid media on spore germination	71
	Washing spores and increasing humidity on germination rates	71
	Using microcycle conidiation versus normal on germination rates	72
	Concentrations of inoculum and broth on germination rates	72
	Droplet Test Development	73
	Controls	73
	Optimized Test	74

Validation bioassay : Sensitivity of <i>Colletotrichum gloeosporioides</i> to Azoxystrobin	74
Baseline bioassay : Sensitivity of <i>Colletotrichum gloeosporioides</i> to Azoxystrobin	78
Validation bioassay : Sensitivity of <i>Colletotrichum capsici</i> to Azoxystrobin	79
Baseline bioassay : Sensitivity of <i>Colletotrichum capsici</i> to Azoxystrobin	83
Summary of Data Analysis	84
Discussion	90
V MOLECULAR MARKER STUDIES	
Introduction	97
Material and Methods	97
Sample Preparation	97
Genomic DNA Extraction	98
Phenol Chloroform Extraction	99
CTAB Extraction Method	100
DNA Purity and Quantity	100
Random Amplified Polymorphic DNA (RAPD) Procedure	101
Long-Primer Random Amplified Polymorphic DNA (LP-RAPD) Procedure	102
Random Amplified Microsatellites (RAMS) Procedure	103
Statistical Analysis of Random Amplified Polymorphic DNA, Long-Primer Random Amplified Polymorphic DNA, Random Amplified Microsatellites markers	105
Results	106
Optimization and the Effect of Different Reaction Parameters	106
Screening and Selection of Random Amplified Polymorphic DNA Primers	106
Random Amplified Polymorphic DNA Bands	108
Band Patterns	109
Random Amplified Polymorphic DNA Population Analysis	109
Screening for Selection of Long Primer - Random Amplified Polymorphic DNA Analysis	109
Long Primer - Random Amplified Polymorphic DNA Bands	119
Screening and Selection of Random Amplified Microsatellites Primers	119
Random Amplified Microsatellites Bands	119
Random Amplified Polymorphic DNA on <i>C. gloeosporioides</i>	124
Random Amplified Polymorphic DNA on <i>C. capsici</i>	124
Long Primer - Random Amplified Polymorphic DNA Trees	127
Long Primer - Random Amplified Polymorphic DNA on <i>C. gloeosporioides</i>	127

Long Primer - Random Amplified Polymorphic DNA on <i>C. capsici</i>	127
Random Amplified Microsatellites Trees	130
Random Amplified Microsatellites on <i>C. gloeosporioides</i>	130
Random Amplified Microsatellites on <i>C. capsici</i>	131
Dendrogram for <i>C. gloeosporioides</i>	131
Dendrogram for <i>C. capsici</i>	131
Discussion	136
VI GENERAL DISCUSSION	146
VII CONCLUSION	156
REFERENCES	158
APPENDICES	
VITAE	

LIST OF TABLES

Tables	Page
3.1 Collection data for <i>Colletotrichum gloeosporioides</i>	47
3.2 Collection data for <i>Colletotrichum capsici</i>	48
3.3 Cultural characteristics of <i>Colletotrichum gloeosporioides</i> from mango grown on Potato Dextrose Agar	51
3.4 Cultural characteristics of <i>Colletotrichum capsici</i> from chilli grown on Potato Dextrose Agar	53
4.1 Amount of Azoxystrobin incorporated into agar for various final concentrations	63
4.2 Germination rates for <i>Colletotrichum gloeosporioides</i> and <i>Colletotrichum capsici</i>	71
4.3 Table of ED ₈₀ estimates and their 95% confidence limits for validation <i>Colletotrichum gloeosporioides</i> isolates	77
4.4 Table of confidence limits for ED ₈₀ values based on validation study	81
4.5 Table of ED ₈₀ estimates and their 95% confidence limits for validation <i>C. capsici</i> isolates.	82
4.6 Validation test results for <i>Colletotrichum gloeosporioides</i>	86
4.7 Validation test results for <i>Colletotrichum capsici</i>	86
4.8 Calculation of confidence limits (95%) around maximum and minimum sensitivity value for <i>Colletotrichum gloeosporioides</i> and <i>Colletotrichum capsici</i>	86
4.9 Azoxystrobin Baseline sensitivity values <i>Colletotrichum gloeosporioides</i>	87
4.10 Azoxystrobin Baseline sensitivity values <i>Colletotrichum capsici</i>	88
5.1 List of RAPD primers used in the population studies	107
5.2 Band frequencies of 22 RAPD zones for 23 individuals representing 5 populations of <i>Colletotrichum gloeosporioides</i>	110
5.3 Band frequencies of 17 RAPD zones for 23 individuals representing 5 populations of <i>Colletotrichum capsici</i>	112

5.4	List of LP-RAPD primers used in the population studies	114
5.5	Band frequencies of 18 LP-RAPD zones for 23 individuals representing 5 populations of <i>Colletotrichum gloeosporioides</i>	115
5.6	Band frequencies of 13 LP-RAPD zones for 23 individuals representing 5 populations of <i>Colletotrichum capsici</i>	117
5.7	List of RAMS primers used in the population studies	118
5.8	Band frequencies of 19 RAMS zones for 23 individuals representing 5 populations of <i>Colletotrichum gloeosporioides</i>	120
5.9	Band frequencies of 18 RAMS zones for 23 individuals representing 5 populations of <i>Colletotrichum capsici</i>	122

LIST OF FIGURES

Figure	Page
3.1 Map of collection sites for <i>Colletotrichum gloeosporioides</i> and <i>Colletotrichum capsici</i> from Malaysia	39
4.1 Alkyl Ester Broth (2x)	60
4.2 Layout of 6 well plates	62
4.3 Dilution series for Azoxystrobin incorporated into agar for various final concentrations	62
4.4 Dilution series for Azoxystrobin rates used	66
4.5 Frequency distribution of Geometric Mean ED ₈₀ values for <i>Colletotrichum gloeosporioides</i>	76
4.6 Histogram of log ED ₈₀ estimates for <i>Colletotrichum gloeosporioides</i> isolates	80
4.7 Graph of log ED ₈₀ estimates for validation <i>C. capsici</i> isolates	81
4.8 Histogram of log ED ₈₀ estimates for <i>Colletotrichum capsici</i> isolates	84
4.9 Frequency distribution of geometric Mean ED ₈₀ values for <i>Colletotrichum gloeosporioides</i>	89
4.10 Frequency distribution of geometric Mean ED ₈₀ values for <i>Colletotrichum capsici</i>	89
5.1 Table of isolates used and phenetic tree for <i>C. gloeosporioides</i> isolates screened with RAPD primers.	125
5.2 Table of isolates used and phenetic tree for <i>C. capsici</i> isolates screened with RAPD primers.	126
5.3 Table of isolates used and phenetic tree for <i>C. gloeosporioides</i> isolates screened with LP-RAPD primers.	128
5.4 Table of isolates used and phenetic tree for <i>C. capsici</i> isolates screened with LP-RAPD primers	129
5.5 Table of isolates used and phenetic tree for <i>C. gloeosporioides</i> isolates screened with RAMS primers	132

5.6	Table of isolates used and phenetic tree for <i>C. capsici</i> isolates screened with RAMS primers	132
5.7	Dendrogram for <i>Colletotrichum gloeosporioides</i>	134
5.8	Dendrogram for <i>Colletotrichum capsici</i>	135

LIST OF ABBREVIATIONS

Chemicals

CTAB	cetyltrimethylammonium bromide
EDTA	Ethylenediamine tetraacetate
KCl	Potassium chloride
MgCl ₂	Magnesium chloride
SDS	Sodium dodecyl sulphate
TBE	Tris-borate-EDTA buffer
TE	Tris-EDTA buffer

Media

AEA	Alkyl Ester Agar
AEB	Alkyl Ester Broth
CB	Cooke's Broth
CDA	Czapek Dox Agar
CMA	Corn Meal Agar
DWA	Distilled Water Agar
LBA	Lima Bean Agar
MA	Malt Agar
MEA	Malt Extract Agar
MEB	Malt Extract Broth
OMA	Oatmeal Agar
PDA	Potato Dextrose Agar
PDB	Potato Dextrose Broth
SDA	Sabourand Dextrose Agar
TWA	Tap Water Agar

Units

°C	Celsius
ED	Effective Dose
hr	hour
kb	kilobase
min	minute
ml	milliliter
mM	millimolar
mg	milligram
nmole	nanomole
ppm	part per million
p.s.i.	pound per square inch
sec	second
spores/ml	spores per milliliter of spore suspension
µg	microgram
µl	microliter
µm	micrometre
µM	micromolar
V	volt
w/v	weight / volume

CHAPTER 1

INTRODUCTION

The disease anthracnose is a prevalent post-harvest disease in South East Asia. It is caused by several fungi, the most common of which is *Colletotrichum*. Certain species of the genus are known to have a wide host range while others have specific hosts. Among members of the genus, the most common form is *Colletotrichum gloeosporioides*, which has a wide host range. It causes symptoms of black sunken lesions on surfaces of fruits such as mangoes, papayas and apples (Freeman *et al.*, 1998) which reduces the market value of the otherwise perfect fruits. An equally important species is *C. capsici* which has a narrow host range, infecting only chillies (Misra and Mahmood, 1960). Mangoes and chillies are the major hosts for *C. gloeosporioides* and *C. capsici* respectively. If not controlled, anthracnose is capable of causing huge post-harvest losses. The most common means of control throughout the world is by the use of fungicides.

The use of fungicide depends on the expectation of achieving a yield greater than its input and its application costs. Compared to other methods of increasing yield, fungicides are comparatively more effective and economical in the long term. Fungicide-treated plants showed significantly greater yield in weight of marketable fruits (Torgeson, 1967).

A fungicide is a molecule that is able to destroy or halt the development of fungi without killing the host (Torgeson, 1967). In the simplest term, a fungicide is a substance destructive to fungi (Abercrombie *et. al.* 1971). There are two general types of fungicides, namely protectants and systemic. Protectants act by shielding the plant surface whereas systemic fungicides are absorbed and translocated to growing points of the plants. Eradicants are used to control disease spread on plants with pronounced diseases symptoms. Fumigants are volatile compounds that protects the whole plant. Currently, classical protective organic fungicides are commonly used, although the usage of highly active systemic compounds with curative action are increasingly being tested (Torgeson, 1967).

Of the agrochemicals used worldwide, fungicides make up only 19% compared to herbicides and insecticides, which makes up 48% and 28% of total pesticide sales in 1997 respectively. Global sales of pesticides amounted to US \$31.25 billion in 1997 (Anon, 1997). Adverse trading conditions during the economic downturn of 1998-1999 have caused a decrease in pesticide sales volume of 5-10%. However, the converse was true for Azoxystrobin where sales grew over 100% after its registration in the United Kingdom, Europe and United States of America in 1997. In 1999, world wide sale of US \$415 million was achieved making Azoxystrobin a leading proprietary fungicide in the world (Anon, 1999).

Azoxystrobin, Zeneca Agrochemical's new broad spectrum systemic fungicide from the strobilurin group, originated from natural fungicides, namely oudemansin A and strobilurin A that are produced by a number of edible mushrooms that grow in temperate forests on decaying wood. These natural anti-fungal products helped the

mushroom to compete with other fungi for nutrients. The compounds were tested and found to give promising results of disease control. This led to a programme of chemical synthesis to design analogues with improved activity and optimized physical properties, such as greater photostability and low volatility. Further research, led to the selection of compounds which showed a high level of fungicidal activity with excellent crop safety in addition to a low mammalian toxicity and benign environmental profile. This compound was identified as Azoxystrobin and was given the registered trade name of Amistar® (Anon, 1996).

Presently, registrations have already been granted in over 40 countries. In the USA, 'Amistar' has been registered under the EPA's Reduced Risk Pesticide Program. In the European Community, it is the first new active ingredient to be Annex I listed. Registrations have been granted or are pending in several Asian countries with widespread sales anticipated in the region from 1999 onwards (Dale *et al.*, 1999).

In March 2000, Azoxystrobin (Amistar®) was awarded the 'Millennium Product' status by the Design Council of United Kingdom for being a product that challenges existing conventions, is environmentally responsible, and demonstrates the application of a new technology. The product has been approved for use in 61 countries on more than 65 crops (Anon, 2000).

In situations of intensive usage it is important to have appropriate monitoring to detect shifts in sensitivity before the product becomes commercially significant. The basis of monitoring is to ensure product effectiveness with prolonged use. In order to do this, the sensitivity of the pathogenic fungi to Azoxystrobin before product